



PREVALENCE OF PRE-DIABETES OR DIABETES AMONG TUBERCULOSIS PATIENTS IN NORTHEAST OF CHINA

Mubinjonov Aziz Daminovich

Jarkinova Marjona G'olibjon qizi

Norqulov Mehrojiddin Mahammad o'g'li

Samarkand State Medical University

Uzbekistan. Samarkand

<https://doi.org/10.5281/zenodo.13734325>

Annotation: tuberculosis (TB) and diabetes are both severe chronic diseases. They could begin simultaneously and also aggravate each other. That is why it is very significant to explore the interaction between two diseases. The goal of this study is to investigate the prevalence rate of TB patients combined with pre-diabetes (impaired fasting blood glucose) or diabetes. The data show that the prevalence rate of TB patients suffering from pre-diabetes or diabetes is 12,4 % and 13,3 %, respectively, which are higher than in general population. These results indicate that TB may play an important role in inducing or promoting the development of diabetes. Therefore, further evidence of the connection between TB and diabetes may provide a new strategy of prevention and treatment of both diseases.

Key words: TB, pre-diabetes, diabetes, combination.

Recent data for the global burden of disease reflect major demographic and lifestyle changes, leading to a rise in non-communicable diseases. Most countries with high levels of tuberculosis (TB) face a large comorbidity burden from both non-communicable and communicable diseases. As the World Health Organization (WHO) and the International Organization against TB and Lung Disease raised a cooperation framework in 2011, which proposed merger care and disease control in diabetic patients with TB, encouraging collaborative research and bidirectional screening for two diseases, the relationship between diabetes and TB or synergies caused more and more concern [1, 2]. Both diabetes and TB are severe chronic diseases. They could outset simultaneously and motivate each other as well. The traditional specific therapeutic methods have failed in management and control of these diseases [3]. Most studies have proposed that diabetes is one of the important risk factors for TB. Diabetes can not only lead to more severe mycobacterium infection, high mortality and low cure rate, but also can lead to a recurrence and reproduce of TB [4, 5]. However, there is less research on TB merger pre-diabetes and diabetes, especially the lack of effective methods in disease control and treatment. Therefore, concerted



therapy method is urgently needs for this situation. The need for sustained and increased funding for these initiatives is greater than ever and requires increased research commitment. The prevalence of pre-diabetes and diabetes is 9,7 % and 15,5 %, respectively in China [6]. Among 3017 people in Nangang district of Harbin in Heilongjiang province, the prevalence for pre-diabetes is 3,75 %, and for diabetes is 10,21 % [7]. All data above is the study of natural populations. According to the diagnostic criteria for diabetes in China (1999 WHO diagnostic criteria), all TB patients will be classified as TB patients, TB patients combined pre-diabetes and TB patients combined diabetes. We investigate the prevalence of TB combined with prediabetes or diabetes, providing the theoretical basis of TB and diabetes interaction, which would be helpful to control TB and reduce mortality.

Materials and methods Study Population To analyze the incidence of TB in Heilongjiang province, we investigated the cases of TB patients from 2001–2011, the data of which is from Heilongjiang province hospital. Random collected 421 clinical TB cases of March ~ May 2011 from Heilongjiang province hospital, and weed out the cases below normal blood glucose levels, then 404 cases are included in this study. According to the diagnostic criteria for diabetes in China (1999 WHO diagnostic criteria), all TB patients will be classified as three group, single TB patients group with blood glucose 3,9~6,1 mmol/L, TB patients combined prediabetes group with blood glucose 6,1~7,0 mmol/L, and TB patients combined diabetes group with blood glucose more than 7,0 mmol/L. Statistical Analysis SPSS16.0 statistical software is applied to processed all data. The prevalence of TB patients combined prediabetes or diabetes are analyzed with 95 % confidence intervals (95 % CI). Ethical Support All information collected in this study have approved by infectious hospital of Heilongjiang province and Harbin Medical University ethics committee.

Results Case numbers of TB and pulmonary TB in Heilongjiang province To analyze the prevalence of TB in Heilongjiang province, we first collect TB cases from 2001–2011 in Heilongjiang province hospital. As shown in Figure 1, the numbers of TB cases increase gradually since 2001, and fall slightly at 2011, but still at a high level. Pulmonary TB is the most important composition of TB, and its incidence trend is the similar to TB, also raise gradually from 2001~2011. Total 404 TB patients cases, 264 cases of male and 140 cases of female, are collected in this study, with an average age of $42,02 \pm 17,518$. According to the fasting blood glucose levels, all TB patients will be classified as three group, single TB patients group with blood glucose 3,9~6,1 mmol/L, TB patients



combined pre-diabetes group with blood glucose 6,1~7,0 mmol/L, and TB patients combined diabetes group with blood glucose more than 7,0 mmol/L. We investigate the prevalence of TB combined with pre-diabetes or diabetes, and the results are shown in Table 1. The morbidity of TB patients with pre-diabetes is 12.4% (95 % CI 7,3%-17,5 %), and the morbidity of TB patients with diabetes is 13,3 % (95 % CI 7,9 %-18,7 %). Table Prevalence of TB patients combined pre-diabetes or diabetes Group (n=404) Case number Morbidity 95 % CI TB 296 TB+Pre-D 50 12,4 % 7,3 %-17,5 % TB+D 58 14,4 % 7,9 %-18,7 % Notes: TB represents single TB patients, TB+Pre-D represents TB patients combined pre-diabetes, and TB+D represents TB patients combined diabetes.

Discussion TB is a major public health threat in the world, nearly a third people infected by mycobacterium TB, and the numbers still develop with a 9,4 million increase every year. When confronted with TB, a chronic and long term illness, households face not only a physical and mental burden, but also an economic burden, which commonly includes the direct costs of treatment, medicine, informal payments, and the indirect costs of a reduction in their ability to generate income [8], especially in low- and middle-income countries. The prevention of TB relies mainly on early identification, prevention of latent TB infection reactivation, and infection control measures [9]. In this study, the data show that prevalence of TB patients in Heilongjiang province increase gradually from 2001–2011, so the trend of Pulmonary TB, the most important composition of TB. Diabetes is also a chronic disease with a develop people suffering from it. The prevalence of diabetes usually occur with territoriality due to development of economic, custom of living or eating. The rising prevalence of diabetes thus poses an increasing risk for the control and prevention of TB [10]. As such, while this case study is focused on a single city in a developing health care system. TB and diabetes have strong co-morbidities, with both the incidence and severity of being affected by each other [11, 12], the causes are still not completely known. We performed this population-based study to investigate morbidity rates of pre-diabetes or diabetes in TB patients, thus provide a theoretical basis data for further exploring the interaction between TB and diabetes. In this study, the morbidity of TB patients with pre-diabetes is 12,4 % (95 % CI 7,3 %-17,5 %), and the standardized rate is 15,0 %, which is 9,7 % higher than China's domestic normal population in 2010 [6], 3,75 % higher than the community population in Nangang district of Harbin in Heilongjiang province in 2009, suggesting TB facilitate the occurrence of pre-diabetes. Meanwhile, the morbidity of TB patients with diabetes is 14,4 % (95 % CI 7,9-



18,7 %) in our study, and the standardized rate is 12,6 %, within the higher level of normal incidence (2 %-12,9 %) [13, 14]. In addition, this morbidity is also higher than 10,21 % from the community population in Nangang district of Harbin in Heilongjiang province in 2009, further indicating TB facilitate the occurrence of diabetes. Total 255 TB patients, 60,6 % of the whole accounted for survey, are investigated with the age from 24~60 years old. Among them, there are 106 TB patients combined prediabetes and diabetes, and 73 patients are the age of 24~60 years old, which is a high percentage (67,6 %) of total cases in study. Therefore, we should focus on the publicity and education of synergy diseases for the patients aged from 24~60 years old, further to decrease the morbidity of synergy diseases. Currently, new therapy is urgently need to treat the combination diseases between TB and diabetes, and people have tried many methods. It is said that application of IFNs play a supporting role for treatment of patients with refractory multiple drug-resistant TB combined diabetes, besides antibiotic treatment [15]. And our study preliminarily analyze the links between TB and diabetes, this association may provide a new strategy concerning the prevention and treatment of both diseases. In conclusion, our study suggest that the prevalence rate of TB patients suffering from pre-diabetes or diabetes is higher than general people.

References:

1. Супхонов, У. У., Файзиев, Х. Ф., Азимова, А. А., & Абдурахмонов, Д. Ш. (2024). ПРОБЛЕМЫ И РЕШЕНИЕ КОНТОРНОЙ ПЛАСТИКИ (ОБЗОРНАЯ ЛИТЕРАТУРА). NAZARIY VA AMALIY FANLARDAGI USTUVOR ISLOHOTLAR VA ZAMONAVIY TA'LIMNING INNOVATSION YO'NALISHLARI, 1(2), 6-17.
2. Супхонов, У. У., Файзиев, Х. Ф., Азимова, А. А., & Абдурахмонов, Д. Ш. (2024). РАССМОТРИМ СРАВНИТЕЛЬНЫЕ АСПЕКТЫ ИСПОЛЬЗОВАНИЯ И ПОБОЧНЫЕ ЭФФЕКТЫ РАЗЛИЧНЫХ МЕТОДОВ ЛИПОАСПИРАЦИИ. NAZARIY VA AMALIY FANLARDAGI USTUVOR ISLOHOTLAR VA ZAMONAVIY TA'LIMNING INNOVATSION YO'NALISHLARI, 1(2), 23-35.
3. Супхонов, У. У., Файзиев, Х. Ф., Азимова, А. А., & Абдурахмонов, Д. Ш. (2024). СУЩЕСТВУЮТ СОВРЕМЕННЫЕ МЕТОДЫ ЛИПОСАКЦИИ, КОТОРЫЕ УСПЕШНО ПРИМЕНЯЮТСЯ ДЛЯ КОНТУРНОЙ ПЛАСТИКИ ТЕЛА. NAZARIY VA AMALIY FANLARDAGI USTUVOR ISLOHOTLAR VA ZAMONAVIY TA'LIMNING INNOVATSION YO'NALISHLARI, 1(2), 18-22.
4. Азимова, А. А., Маликов, Д. И., & Шайкулов, Х. Ш. (2021). МОНИТИРОИНГ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ СЕПСИСА ЗА. PEDAGOGICAL SCIENCES AND TEACHING METHODS, 48.



5. Maksudov, D., Azimova, A., & Mamadyarova, S. (2021). Clinical course of inflammatory diseases of the maxillofacial region in patients with viral hepatitis b. Збірник наукових праць Л'ОГОС.–2021.
6. АЗИМОВА, А. А., & МАЛИКОВ, Д. И. (2022). Анатомические Компоненты Улыбки И Ее Патологии. МОЛОДЕЖНЫЙ ИННОВАЦИОННЫЙ ВЕСТНИК Учредители: Воронежский государственный медицинский университет имени НН Бурденко, 11(S1), 255-257.

